

# Electrogenerated Chemiluminescence. 37. Aqueous Ecl Systems Based on Ru(2,2'-bipyridine)<sub>3</sub><sup>2+</sup> and Oxalate or Organic Acids

Israel Rubinstein and Allen J. Bard\*


Contribution from the Department of Chemistry, The University of Texas at Austin, Austin, Texas 78712. Received June 9, 1980

**Abstract:** An aqueous system for electrogenerated chemiluminescence (ecl) based on the reaction of electrogenerated Ru(bpy)<sub>3</sub><sup>3+</sup> with strong reductants produced as intermediates in the oxidation of oxalate ion is described. The bright orange chemiluminescence, which could also be generated by reaction of chemically produced Ru(III) species with oxalate, corresponded to emission by Ru(bpy)<sub>3</sub><sup>2+\*</sup>; ecl efficiency (photons emitted/Ru(bpy)<sub>3</sub><sup>3+</sup> generated) was ~2% in deaerated solution. Ecl by reaction of the 1+ and 3+ Ru species could also be obtained in partially aqueous solutions containing at least 20% acetonitrile. Chemiluminescence was also observed with other organic acids (pyruvic, malonic, lactic), when the intermediates produced on their oxidation by Ce<sup>4+</sup> reacted with Ru(bpy)<sub>3</sub><sup>3+</sup>.

## Introduction

The numerous investigations of chemiluminescence arising from excited-state production in energetic electron-transfer reactions of electrogenerated reactants (electrogenerated chemiluminescence or ecl) have almost always involved nonaqueous solvent media such as *N,N*-dimethylformamide or acetonitrile (MeCN).<sup>1</sup> Indeed even small amounts of water will frequently quench the ecl reaction and studies are usually carried out under rigorously dry conditions utilizing vacuum line or drybox techniques.<sup>2</sup> The major difficulties with designing ecl systems in aqueous solutions are (a) the very low solubility of the organic compounds which are the radical ion precursors, (b) the reactivity of the radical ions with water, and (c) the limited potential range of water. However, aqueous ecl systems would be of interest, especially for the design of practical ecl devices (e.g., displays). In many cases, the lifetime of an ecl device appears to be governed by the presence of trace impurities (e.g., water) in the solvent and supporting electrolyte.

A very good candidate for aqueous ecl is the Ru(bpy)<sub>3</sub><sup>2+</sup> (where bpy = 2,2'-bipyridine) system. This system involves water-soluble species, and previous studies<sup>3</sup> have shown that in MeCN solutions intense ecl results from the reaction of the 3+ and 1+ species.



An investigation of the temperature dependence of the ecl efficiency (photons emitted per redox event) suggested that the production of the excited state by the redox reaction (1) occurs with an efficiency near 100%.<sup>4</sup> In addition to the annihilation mode of ecl with Ru(bpy)<sub>3</sub><sup>3+/+</sup>, chemiluminescence involving reaction of the +3 species with other reducing agents is possible.<sup>5,6</sup> Thus the simultaneous oxidation of Ru(bpy)<sub>3</sub><sup>2+</sup> and oxalate ion in MeCN at a Pt electrode produces light emission. The mechanism proposed for this emission was based on the oxidation of oxalate ion to form the strong reducing agent CO<sub>2</sub><sup>-</sup>, which ultimately yields an electron transfer to Ru(bpy)<sub>3</sub><sup>3+</sup> to produce an excited state.


We report here aqueous ecl systems involving Ru(bpy)<sub>3</sub><sup>2+</sup> and a preliminary study of coupled reactions ("oxidative reductions") of oxalate and several organic acids which were revealed by the ecl investigations.

## Experimental Section


Electrochemical measurements were performed in a simple three-electrode cell, using Pt wire or a glassy carbon disk (Tokai Carbon Co., Tokyo, Japan) working electrodes (geometric area, 0.20 cm<sup>2</sup>), a Pt auxiliary electrode, and either a mercury sulfate reference electrode [(MSE) Hg/Hg<sub>2</sub>SO<sub>4</sub>/1.0 M H<sub>2</sub>SO<sub>4</sub>; +0.67 V vs. NHE<sup>7</sup>] or a silver wire quasi reference electrode.<sup>8</sup> The electrochemical and ecl experiments generally used the instrumentation and techniques previously described.<sup>3,5</sup> Ecl and fluorescence spectra were measured on an Aminco-Bowman spectrophotofluorometer. Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub> was prepared from the chloride (G. F. Smith Chemical Co.).<sup>3b</sup> Oxalic acid (Mallinckrodt, A.R.), sodium oxalate (Allied Chemical), sodium sulfate (Fisher Scientific Co., certified ACS), cerium(III) sulfate (Alfa Products, 99.9%), ceric ammonium sulfate (G. F. Smith Chemical Co.), lead dioxide (Allied Chemical), tetramethylammonium chloride (Southwestern Analytical Chemicals, polargraphic grade), and acetonitrile (MCB, spectroquality) were all used without further purification. All solutions were prepared with triple-distilled water.

## Results and Discussion

**(A) The Ru(bpy)<sub>3</sub><sup>2+</sup>-Oxalate System. Electrochemical measurements.** Current-potential curves for the Ru(bpy)<sub>3</sub><sup>3+/2+</sup> reversible couple and the anodic oxidation of oxalate at a Pt electrode in 1.0 M H<sub>2</sub>SO<sub>4</sub> are presented in Figure 1a-c. Oxalate is irreversibly oxidized to CO<sub>2</sub> at Pt at potentials within the platinum oxide region. Although this reaction has been investigated before<sup>9</sup> and the only product in aqueous solution is CO<sub>2</sub>, the detailed mechanism is uncertain. The oxalate oxidation wave is known to be completely suppressed on an oxide-covered Pt surface,<sup>9</sup> as is also demonstrated by the voltammogram in Figure 1c. The electroactive species is usually assumed to be the adsorbed undissociated acid, which is oxidized according to the overall eq 3.



In the simultaneous oxidation of oxalate and Ru(bpy)<sub>3</sub><sup>2+</sup>, in 1.0 M H<sub>2</sub>SO<sub>4</sub> at a reduced Pt electrode (Figure 1d), the oxidation peaks of both species occur at nearly the same potential. This results in a single broad oxidation peak. On the other hand, with a Pt electrode which has been preoxidized (an oxide-covered electrode), one observes only the reversible Ru(bpy)<sub>3</sub><sup>3+/2+</sup> couple



Carbon electrodes (both pyrolytic graphite and glassy carbon) show behavior similar to that of oxide-covered Pt electrodes toward

(1) See, for example: (a) Faulkner, L. R. *Methods Enzymology* 1978, 57; (b) Faulkner, L. R.; Bard, A. J. *Electroanal. Chem.* 1977, 10.

(2) An exception is the ecl in the luminol-oxygen system by: Epstein, B.; Kuwana, T. *Photochem. Photobiol.* 1965, 4, 1157; *Ibid.* 1967, 6, 605.

(3) (a) Tokel, N. E.; Bard, A. J. *J. Am. Chem. Soc.* 1972, 94, 2862. (b) Tokel-Takvoryan, N. E.; Hemingway, R. E.; Bard, A. J. *Ibid.* 1973, 95, 6582.

(4) (a) Wallace, W. L.; Bard, A. J. *J. Phys. Chem.* 1979, 83, 1350. (b) Itoh, K.; Honda, K. *Chem. Lett.* 1979, 99.


(5) Chang, M.; Saji, T.; Bard, A. J. *J. Am. Chem. Soc.* 1977, 99, 5399.

(6) Lytle, F. E.; Hercules, D. M. *Photochem. Photobiol.* 1971, 13, 123.

(7) Reichman, B.; Bard, A. J. *J. Electrochem. Soc.* 1979, 126, 583.

(8) Keszthelyi, C. P.; Tachikawa, H.; Bard, A. J. *J. Am. Chem. Soc.* 1972, 94, 1522.

(9) (a) Lingane, J. J. *J. Electroanal. Chem.* 1959/1960, 1, 379. (b) Giner, J. *Electrochim. Acta* 1961, 4, 42. (c) Anson, F. C.; Schulz, F. A. *Anal. Chem.* 1963, 35, 1115. (d) Johnson, J. W.; Wroblowa, H.; Bockris, J. O'M. *Electrochim. Acta* 1964, 9, 639.



**Figure 1.** Current-potential curves at steady state for a Pt electrode (sweep rate, 200 mV/s): (a) 1.0 M H<sub>2</sub>SO<sub>4</sub>; (b) 1.0 mM Ru(bpy)<sub>3</sub><sup>-</sup>(ClO<sub>4</sub>)<sub>2</sub>, 1.0 M H<sub>2</sub>SO<sub>4</sub>; (c) 3.0 mM H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>, 1.0 M H<sub>2</sub>SO<sub>4</sub>; (d) 3.0 mM H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>, 1.0 mM Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub>, 1.0 M H<sub>2</sub>SO<sub>4</sub>; (e) Same as d but 0.1 M H<sub>2</sub>SO<sub>4</sub>; (f) same as d at pH 4.5 (adjusted by addition of NaOH to 0.1 M H<sub>2</sub>SO<sub>4</sub>).


oxalate and Ru(bpy)<sub>3</sub><sup>2+</sup> oxidation; i.e., there is a very large overpotential for oxalate oxidation on carbon compared to that on reduced Pt. The potential at which oxalate oxidation occurs depends upon the pH and becomes less positive with an increase in pH. Thus in 0.1 M H<sub>2</sub>SO<sub>4</sub> (pH ~1.8) the cyclic voltammogram of a solution containing both oxalate and Ru(bpy)<sub>3</sub><sup>2+</sup> shows separated waves (Figure 1e). A further increase in the pH causes an even greater separation (Figure 1f). Moreover, at an oxidized electrode in the presence of oxalate, the anodic current for Ru(bpy)<sub>3</sub><sup>2+</sup> was larger and the cathodic current on the reverse scan was smaller than in its absence (Figure 1e). This results from the reaction of the 3+ species with oxalate, which under these conditions is not oxidized directly at the electrode.

While the mechanism for the oxidation of oxalate by Ru(bpy)<sub>3</sub><sup>3+</sup> or at an electrode has not been established, the previous ecl results<sup>5</sup> and the fact that Ru(III) is usually a one-electron oxidant suggest reaction 5 followed by rapid decomposition of C<sub>2</sub>O<sub>4</sub><sup>-</sup>. (For oxalic




acid, pK<sub>a1</sub> = 1.23 and pK<sub>a2</sub> = 4.19,<sup>10</sup> so that the predominant forms in acidic solution are H<sub>2</sub>C<sub>2</sub>O<sub>4</sub> and HC<sub>2</sub>O<sub>4</sub><sup>-</sup>.) Thus at an oxidized Pt electrode oxalate is oxidized exclusively by reaction with Ru(bpy)<sub>3</sub><sup>3+</sup>, thus producing in the diffusion layer near the electrode a mixture of the 2+ and 3+ Ru species as well as intermediates from the oxidation of oxalate. The catalytic oxidation of oxalate by Ru(bpy)<sub>3</sub><sup>3+</sup> does not occur in 1 M H<sub>2</sub>SO<sub>4</sub>, however.

(10) Weast, W. C. "Handbook of Chemistry and Physics", 51st ed.; CRC Press: Cleveland, Ohio, 1970; p D-120.



**Figure 2.** (a) Ecl emission spectrum of Pt electrode in a solution of 1.0 mM Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub> and 50 mM Na<sub>2</sub>C<sub>2</sub>O<sub>4</sub> at pH 5.0, with cyclic square wave excitation, between 0.2 and 0.7 V vs. MSE at 0.2 Hz. (b) Luminescence emission spectrum of 1.0 mM Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub> in 0.1 M H<sub>2</sub>SO<sub>4</sub>, with photoexcitation at 500 nm.



**Figure 3.** Dependence of the ecl intensity (at 610 nm) on the pH. The solution contained 1.0 mM Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub> and 6.0 mM H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>, with the pH adjusted by addition of NaOH to 0.1 M H<sub>2</sub>SO<sub>4</sub>.


**Electrogenerated Chemiluminescence.** When a potential which is more positive than the potential for the oxidation of Ru(bpy)<sub>3</sub><sup>2+</sup> is applied to a Pt or carbon electrode in an aqueous solution containing as low as 1.0 mM Ru(bpy)<sub>3</sub><sup>2+</sup> and various concentrations of oxalate at pH ≥ 2, orange light is emitted from the electrode surface. No ecl occurs in 1.0 M H<sub>2</sub>SO<sub>4</sub>. The ecl can be produced either by repetitively pulsing the electrode potential between a potential sufficiently positive for Ru(bpy)<sub>3</sub><sup>2+</sup> oxidation and a more negative rest potential to allow renewal of the depleted oxalate near the electrode surface or by applying a constant positive potential with stirring of the solution. With high oxalate concentration stirring is unnecessary. The ecl intensity is quite strong (clearly visible to the non-dark-adapted eye) even at low concentrations (e.g., 5 mM) of oxalate at 7 ≥ pH ≥ 4. Removal of oxygen from the solution is not necessary, but deaeration with nitrogen for several minutes increases the ecl intensity. This light emission is clearly attributable to the excited state, Ru(bpy)<sub>3</sub><sup>2+\*</sup>, since the ecl and luminescence spectra are essentially identical (Figure 2).

No emission is observed at potentials where only oxidation of oxalate occurs; potentials sufficiently positive to cause Ru(bpy)<sub>3</sub><sup>2+</sup> oxidation are required. In addition, in the repetitive pulsing mode, a shift in the rest potential to sufficiently negative values to cause the removal of the oxide layer (which is present on the Pt surface at the potential for oxidation of Ru(bpy)<sub>3</sub><sup>2+</sup>) results in a substantial decrease in the emission intensity. The ecl intensity is strongly dependent on the pH, as shown in Figure 3. Note that these measurements were made in unbuffered solutions, so that the pH at the electrode surface during oxidation was somewhat lower than


in the bulk solution. Addition of buffers, e.g., acetate, decreased the ecl intensity. However, over a pH range of 4–6 with an acetate buffer of the same total acetate concentration the trend in the variation of ecl intensity with pH was qualitatively the same as in the unbuffered solutions.

The ecl efficiency was estimated by comparing the ratio of the integrated light intensity to electric charge passed for the present system to that of the Ru(bpy)<sub>3</sub><sup>2+</sup>–acetonitrile nonaqueous system, whose ecl efficiency is known.<sup>4</sup> In an aqueous deaerated solution of 1.0 mM Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub> and 50 mM Na<sub>2</sub>C<sub>2</sub>O<sub>4</sub>, the  $\phi_{\text{ecl}}$  calculated in this manner, taking into account that each ecl event requires the transfer of two electrons, was ~2%. In a non-deaerated solution,  $\phi_{\text{ecl}}$  decreased to about one-third of this value.

The mechanism of this ecl reaction clearly involves reaction of Ru(bpy)<sub>3</sub><sup>3+</sup> and an intermediate formed on oxidation of oxalate. The excited state cannot be produced via reaction 5 since the free energies of the Ru(bpy)<sub>3</sub><sup>3+/2+</sup> and C<sub>2</sub>O<sub>4</sub><sup>2-</sup>/CO<sub>2</sub> couples are insufficient.<sup>11</sup> At the applied electrode potential for the ecl generation (more positive than that of the Ru(bpy)<sub>3</sub><sup>3+/2+</sup> couple), the Pt electrode is covered with an oxide layer, which prevents the direct oxidation of oxalate on the electrode (see Figure 1c). On the other hand, at pH values above ~3, the generated Ru(bpy)<sub>3</sub><sup>3+</sup> oxidizes the oxalate (as shown in Figure 1e) near the electrode, according to (5). This can be followed by



The intermediate radical CO<sub>2</sub><sup>·-</sup>, which results from the one-electron oxidation of oxalate, is a strong reducing agent and can produce the excited state by reacting directly with the 3+ species



followed by (2) or possibly by



followed by (1) and (2).

We consider here both (7) and (8) as possible routes for generation of the excited state, although (8) was shown to be more favorable in MeCN.<sup>5</sup>

At lower pHs where the predominant species in solution is HC<sub>2</sub>O<sub>4</sub><sup>-</sup>, one would expect a reaction sequence similar to (5)–(8) with the reactive intermediate now being CO<sub>2</sub>H<sup>·</sup>, which would be a weaker reducing agent than CO<sub>2</sub><sup>·-</sup>. This will account for the pH dependence shown in Figure 3, where the increase in the ecl intensity occurs in the vicinity of pK<sub>a2</sub> of the oxalic acid and roughly follows the fractional concentration of C<sub>2</sub>O<sub>4</sub><sup>2-</sup> in solution. Another factor which may contribute to the increase in intensity with the pH is the shift of the potential for oxidation of oxalate, as shown in Figure 1. The decrease in the ecl intensity above pH 6 is most likely caused by the increasing importance of the oxygen evolution reaction at the electrode, which is shifted to less positive potentials with increasing pH. The increasing importance of the reaction of Ru(bpy)<sub>3</sub><sup>3+</sup> with water (or hydroxide ion) in this pH region<sup>12</sup> may also play a role in this decrease. Very weak emission exists even at pH values above 10; this is probably caused by the known light-emitting reduction of Ru(bpy)<sub>3</sub><sup>3+</sup> with OH<sup>-</sup> ions.<sup>6</sup>

Note that each oxalate species directly oxidized on the electrode is wasted with respect to the ecl. Since CO<sub>2</sub><sup>·-</sup> (or CO<sub>2</sub>H<sup>·</sup>) is a much stronger reducing agent than C<sub>2</sub>O<sub>4</sub><sup>2-</sup> (or HC<sub>2</sub>O<sub>4</sub><sup>-</sup>), when this intermediate is formed at the electrode surface it will immediately transfer the second electron to the electrode to produce CO<sub>2</sub>. The oxide layer on the Pt surface thus prevents the direct oxidation of oxalate at the electrode. This accounts for the experimental observation that the ecl intensity decreases when the oxide layer is repeatedly removed in the pulsed-mode ecl experiments. In this respect, carbon electrodes have the same advantage as oxidized Pt electrodes, since they show a large overpotential for the electrooxidation of oxalate so that it occurs at more positive




Figure 4. Flow system for continuous light generation: a and b, glass compartments; c, magnetic stirrer; d, solid BaC<sub>2</sub>O<sub>4</sub>; e, solid PbO<sub>2</sub>; f, peristaltic pump.



potentials than those for the reversible oxidation of Ru(bpy)<sub>3</sub><sup>2+</sup>.

**Solution Chemiluminescence Reactions.** Light emission from the same system can be generated not only at an electrode surface but also in homogeneous solution by chemical reaction. Quantitative oxidation of an orange solution of Ru(bpy)<sub>3</sub><sup>2+</sup> in 0.1 M H<sub>2</sub>SO<sub>4</sub>, either electrochemically or with PbO<sub>2</sub>, produces a green solution of Ru(bpy)<sub>3</sub><sup>3+</sup>. Addition of sodium oxalate (or oxalic acid) powder to this solution immediately results in a very strong orange luminescence. In agreement with the ecl results, only a very low-intensity emission is generated if the same experiment is performed in 1.0 M H<sub>2</sub>SO<sub>4</sub>. These experiments suggest that the ecl emission is indeed the result of a homogeneous reaction between Ru(bpy)<sub>3</sub><sup>3+</sup> and oxalate.

**Chemiluminescent Systems.** The Ru(bpy)<sub>3</sub><sup>2+</sup>–oxalate aqueous ecl system presents some important advantages for possible practical applications. It is an aqueous, very stable, and intense ecl system, relatively insensitive to the presence of oxygen or organic and inorganic impurities. The main disadvantage is that it is not a regenerative system, since oxalate is consumed. Still, because oxalate does not react directly at an oxidized Pt electrode, one can keep an ecl system operating for prolonged periods of time by having a sufficiently large amount of oxalate in the solution. In a typical experiment, a constant anodic current of 6.8 mA cm<sup>-2</sup> was applied to a Pt flag electrode (area, 0.22 cm<sup>2</sup>) in a solution containing 1.0 mM Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub> and 0.25 M Na<sub>2</sub>C<sub>2</sub>O<sub>4</sub> in acetate buffer (pH 4.8). A strong, constant ecl was observed (without stirring) for 20 h, at which time the experiment was terminated.

Another possibility is to make use of a chemical oxidizing agent in a light-generating device. This was done with the flow system shown in Figure 4. The solution flowing in a closed circuit was 1.0 mM Ru(bpy)<sub>3</sub>Cl<sub>2</sub> in 0.1 M H<sub>2</sub>SO<sub>4</sub>, and the solids in the two compartments were gently stirred. The slightly soluble BaC<sub>2</sub>O<sub>4</sub> in compartment a provides a constant supply of oxalate ions to compartment b, where they react with Ru(bpy)<sub>3</sub><sup>3+</sup> that was produced by oxidation of Ru(bpy)<sub>3</sub><sup>2+</sup> by PbO<sub>2</sub>. As a result, one observes a constant light emission from compartment b. The use of a slightly soluble oxalate and two separate compartments was dictated by the desire to minimize loss of oxalate and PbO<sub>2</sub> by direct reaction between the two species. Here again the light emission lasted until the oxalate and PbO<sub>2</sub> were consumed.


**(B) The Ru(bpy)<sub>3</sub><sup>3+/+</sup> Regenerative System.** Ru(bpy)<sub>3</sub><sup>2+</sup> provides a regenerative ecl system in acetonitrile,<sup>3</sup> when a Pt electrode potential is pulsed between the oxidation and reduction potentials of the complex




followed by (1) and (2). This, however, cannot be achieved in water. Ru(bpy)<sub>3</sub><sup>+</sup> is not generated, even at a glassy carbon electrode, which shows a large overpotential for hydrogen evolution. Figure 5 presents the electrochemical behavior of Ru(bpy)<sub>3</sub><sup>2+</sup> in aqueous solution on glassy carbon electrode. The behavior (on the reduction side) is similar to that with a mercury cathode. On the anodic side, one observes the reversible oxidation

(11) Latimer, W. M. "Oxidation Potentials", Prentice-Hall: New York, 1952.

(12) Creutz, C.; Sutin, N. *Proc. Natl. Acad. Sci. U.S.A.*, 1975, 72, 2858.



**Figure 5.** Current-potential curve at steady state for a glassy carbon disk electrode (sweep rate, 200 mV/s) in a solution containing 1.0 mM Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub> in aqueous 0.1 M Na<sub>2</sub>SO<sub>4</sub>. Dashed curve shows background current in 0.1 M Na<sub>2</sub>SO<sub>4</sub>.




**Figure 6.** Current-potential curve at steady state for a glassy carbon electrode (sweep rate, 200 mV/s) in H<sub>2</sub>O-CH<sub>3</sub>CN (1:1 by volume) containing 1.0 mM Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub> and 0.1 M tetramethylammonium chloride.

of Ru(bpy)<sub>3</sub><sup>2+</sup> (eq 9). On the cathodic side, no reduction wave corresponding to the formation of Ru(bpy)<sub>3</sub><sup>+</sup> is observed. The cathodic peak preceding the catalytic hydrogen evolution is probably a two-electron reduction leading to formation of Ru(bpy)<sub>3</sub><sup>0</sup> accompanied by decomposition and precipitation of Ru<sup>0</sup> species on the electrode. This is evident from the shape of the peaks and also by the observation of decomposition product. If the electrode is held at the cathodic peak potential, the formation of a black precipitate in the solution is seen. No ecl is generated on pulsing the electrode potential between the anodic and cathodic peak potentials.

Upon addition of acetonitrile to the aqueous solution, the electrochemical behavior of the system changes, as shown in Figure 6. A second, more positive reduction peak now appears, which probably corresponds to reaction 10. The height of this Ru(bpy)<sub>3</sub><sup>+</sup> peak gradually becomes smaller with decreasing percentages of acetonitrile in the solution.



Bright ecl is generated upon pulsing the potential of a glassy carbon electrode between the oxidation peak and the first reduction peak of Ru(bpy)<sub>3</sub><sup>2+</sup> in an aqueous solution of 1.0 mM Ru(bpy)<sub>3</sub><sup>2+</sup>, which contains at least 20% acetonitrile. Below 20% acetonitrile no ecl is observed. When the negative potential pulse extends to the second reduction peak region, the light emission decreases or even disappears. This aqueous ecl system is not stable for long periods of operation, however. The light intensity decreases during operation and finally stops, within 5–30 min. This is probably the result of the instability of the 1+ complex toward further



**Figure 7.** Current-potential curves at steady state for a Pt electrode (sweep rate, 200 mV/s) in 0.1 M H<sub>2</sub>SO<sub>4</sub>: (a) 5.0 mM sodium pyruvate and 1.0 mM Ru(bpy)<sub>3</sub>(ClO<sub>4</sub>)<sub>2</sub>; (b) 0.58 mM Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>; (c) 0.58 mM Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> and 5.0 mM sodium pyruvate.

reduction or decomposition in aqueous solution.

**(C) Emission with Other Organic Acids.** Figure 7a presents the electrochemical behavior of the system Ru(bpy)<sub>3</sub><sup>2+</sup>-pyruvate at a Pt electrode in 0.1 M H<sub>2</sub>SO<sub>4</sub>. As in the case of oxalate, the oxidation of pyruvate is suppressed by the presence of an oxide layer on the electrode. The potentials for oxidation of the two species are almost identical on a reduced Pt electrode, and the Ru(bpy)<sub>3</sub><sup>2+</sup> oxidation on oxide-covered Pt is reversible. No ecl is generated upon stepping the potential to values past the peak. All of these observations are similar to the case of Ru(bpy)<sub>3</sub><sup>2+</sup>-oxalate in 1.0 M H<sub>2</sub>SO<sub>4</sub>. The electrochemical behavior of Ce<sup>4+/3+</sup> is shown in Figure 7b. Ce<sup>3+</sup> is reversibly oxidized in 0.1 M H<sub>2</sub>SO<sub>4</sub>; the peaks appear small because the concentration is low. The results of addition of sodium pyruvate to this system are presented in Figure 7c. On the oxidized Pt, the Ce<sup>3+</sup> now produces a much larger anodic current, with a very small reverse peak because of the reaction of Ce<sup>4+</sup> with pyruvate. Since the potential for the oxidation of Ce<sup>3+</sup> is more positive than that for Ru(bpy)<sub>3</sub><sup>2+</sup> oxidation in this system, Ce<sup>4+</sup> is a better oxidant than Ru(bpy)<sub>3</sub><sup>3+</sup> so that reactions 11–13 are now possible.



Addition of Ru(bpy)<sub>3</sub><sup>2+</sup> to this solution and repetitively pulsing to the Ce<sup>3+</sup> oxidation potential result in the generation of the characteristic orange luminescence at the electrode. No ecl is observed on pulsing only to the Ru(bpy)<sub>3</sub><sup>2+</sup> oxidation potential. This suggests that the intermediate radical, CH<sub>3</sub>CO<sup>•</sup>, is a strong enough reducing agent to participate in electron-transfer reactions with the Ru(bpy)<sub>3</sub><sup>3+</sup> similar to (7) or (8). The product of the oxidation of CH<sub>3</sub>CO<sup>•</sup> will be the carbonium ion, which will immediately react with water. This is in agreement with the results on the electrooxidation of pyruvic acid in H<sub>2</sub>O<sup>13</sup> or methanol,<sup>14</sup>

where the main products were reported to be acetic acid and methyl acetate, respectively.

Light emission is also observed when a Ce(IV) salt is added to a solution of Ru(bpy)<sub>3</sub>Cl<sub>2</sub> and sodium pyruvate or when sodium pyruvate is added to a solution of Ru(bpy)<sub>3</sub>Cl<sub>2</sub> containing an excess of Ce<sup>4+</sup>. In both cases, an orange light emission immediately appears. The same chemical experiment generates light emission with lactic and malonic acids and probably will occur with other organic acids or salts that can be rapidly oxidized by Ce<sup>4+</sup>.

Note that one can produce ecl with the Ru(bpy)<sub>3</sub><sup>2+</sup>-oxalate system in 1.0 M H<sub>2</sub>SO<sub>4</sub> by adding Ce<sup>3+</sup> to the solution. The light emission is weak in this case because of the low solubility of Ce(III)-oxalate. A strong emission occurs upon addition of Ce<sup>4+</sup> to a mixture of oxalate and Ru(bpy)<sub>3</sub><sup>2+</sup> in 1.0 M H<sub>2</sub>SO<sub>4</sub> in the chemical experiment. An interesting point is that in the same system (Ru(bpy)<sub>3</sub><sup>2+</sup>, oxalate, 1.0 M H<sub>2</sub>SO<sub>4</sub>) addition of PbO<sub>2</sub> causes only a very low intensity light emission, which develops slowly, in spite of the fact that PbO<sub>2</sub> oxidizes both oxalate and Ru(bpy)<sub>3</sub><sup>2+</sup> very rapidly. This difference in behavior between Ce(IV) and PbO<sub>2</sub> can be explained by the proposed mechanism. Ce<sup>4+</sup>, which is a one-electron oxidant, reacts with oxalate to form the reactive intermediate very efficiently. On the other hand, on the solid PbO<sub>2</sub> the oxalate undergoes a direct two-electron oxidation which is ineffective for light emission. The low level light emitted by adding PbO<sub>2</sub> is probably only the result of the very

slow reaction between Ru(bpy)<sub>3</sub><sup>3+</sup> and oxalate in 1.0 M H<sub>2</sub>SO<sub>4</sub>.

### Conclusions

A new aqueous chemiluminescent and ecl system based on Ru(bpy)<sub>3</sub><sup>2+</sup> and a suitable organic acid or salt (e.g., C<sub>2</sub>O<sub>4</sub><sup>2-</sup>) is described. The system produces quite bright emission and may be useful in practical ecl devices. The system is an example of a coupled chemical reaction<sup>15</sup> with the observation of light emission as a probe of the reaction mechanism. The Ce<sup>3+</sup>-oxalate system (or one with another organic acid or salt) provides a unique electrochemical system for carrying out a reduction under oxidizing conditions at the electrode surface (an "anodic or oxidative reduction"). Upon applying a sufficiently positive potential to an oxidized Pt (or carbon) electrode, one promotes the one-electron oxidation of the acid by the Ce<sup>4+</sup>, creating an intermediate (e.g., CO<sub>2</sub><sup>-</sup>) which is a strong reducing agent. This intermediate has a sufficiently long lifetime in aqueous solution to be able to reduce other molecules in the solution (e.g., Ru(bpy)<sub>3</sub><sup>3+</sup> in the light-emitting reaction in 1.0 M H<sub>2</sub>SO<sub>4</sub>). This system can therefore, in principle, be used for electroreductions in aqueous solution under conditions of mild anodic potential and at a Pt electrode.

**Acknowledgment.** The support of this research by the Army Research Office is gratefully acknowledged.

(14) Wladislaw, B.; Zimmermann, J. P. *J. Chem. Soc. B* 1970, 290.

(15) Laitinen, H. A.; Harris, W. E. "Chemical Analysis"; McGraw-Hill: New York, 1975; pp 297-303.

## Synthesis of "Face to Face" Porphyrin Dimers Linked by 5,15-Substituents: Potential Binuclear Multielectron Redox Catalysts<sup>1a</sup>

James P. Collman,\* Anthony O. Chong, Geoffrey B. Jameson, Richard T. Oakley,<sup>1b</sup> Eric Rose, Eric R. Schmittou,<sup>1c</sup> and James A. Ibers\*

Contribution from the Departments of Chemistry, Stanford University, Stanford, California 94305, and Northwestern University, Evanston, Illinois 60201.  
Received January 25, 1980

**Abstract:** As part of the development of binuclear transition-metal complexes capable of facilitating multielectron redox reactions of small molecules (such as O<sub>2</sub> and N<sub>2</sub>), we have devised an efficient and synthetically flexible route to 5,15-disubstituted porphyrins which affords multigram quantities of isomerically pure porphyrins without recourse to chromatography. The 5,15-substituents are principally amine ((CH<sub>2</sub>)<sub>n</sub>NHR (*n* = 1, 2, 3; R = H, CH<sub>3</sub>)) and acid chloride ((CH<sub>2</sub>)<sub>n</sub>C(O)Cl (*n* = 1, 2)) groups. Reactions of the two types of substituted porphyrins lead to a new class of "face to face" porphyrins with interporphyrin amide bridges of varying length (4-7 atoms). An amine-linked dimer has been produced by the reduction of an appropriate amide linkage. Purely hydrocarbon-linked dimers have also been prepared by the direct coupling of bis(dipyrrylmethanes) bridged by aliphatic (CH<sub>2</sub>)<sub>n</sub> (*n* = 4, 6) chains. The dimeric porphyrin compounds have been fully characterized by their visible and NMR spectra. The crystal and molecular structure of the bis(copper(II)) complex of one of the amide-linked dimers, the first such determination of a "face to face" porphyrin, is reported. Crystal data for Cu<sub>2</sub>C<sub>74</sub>N<sub>10</sub>O<sub>2</sub>H<sub>86</sub>·2H<sub>2</sub>O·C<sub>7</sub>H<sub>8</sub>: space group C<sub>2h</sub>-P2<sub>1</sub>/c; *a* = 11.878 (6) Å, *b* = 13.304 (7) Å, *c* = 23.725 (13) Å, β = 114.60 (2)° at -140 °C; Z = 2. A crystallographic center of symmetry is imposed on the dimer. The structure was refined on all data including F<sub>o</sub><sup>2</sup> < 0 (4663 observations) to values for *R* and *R*<sub>w</sub> on F<sup>2</sup> of 0.103 and 0.152. For the portion of data having F<sub>o</sub><sup>2</sup> > 3σ(F<sub>o</sub><sup>2</sup>) (3213 observations) the values for *R* and *R*<sub>w</sub> on *F* are 0.063 and 0.073. The spectral properties of these new porphyrin dimers are discussed with reference to the structural analysis. A striking feature of the structure is a shear-like displacement of one porphyrin unit with respect to the other by an average distance of 4.95 Å. The Cu...Cu separation is 6.332 (4) Å and the interplanar separation of the two porphyrin rings is 3.87 Å. In solution at ambient temperatures, conformational changes involving motions of the two porphyrin rings cause considerable line broadening in the <sup>1</sup>H NMR spectra of all 5,15-bridged dimers. However, at temperatures above 90 °C conformational equilibria are sufficiently rapid to produce well-resolved spectra.

### Introduction

We have been interested in the development of new catalysts for the direct, four-electron electrochemical reduction of O<sub>2</sub> to H<sub>2</sub>O at potentials approaching the thermodynamic value of +1.23 V (vs. the normal hydrogen electrode). In a broader context such

catalysts may also promote multielectron-transfer reactions for other small molecules such as dinitrogen. In a recent paper we discussed our approach to the design of multielectron redox

\* Address correspondence as follows: J.P.C., Stanford University; J.A.I., Northwestern University.

(1) (a) For a more detailed treatment of materials in this paper see: Schmittou, Eric R. Ph.D. Dissertation, Stanford University, 1979. (b) Natural Sciences and Engineering Council of Canada Postdoctoral Fellow, 1976-1978. (c) National Science Foundation Predoctoral Fellow, 1973-1976.